

Nutritional and Supportive Care

Nutrition During Fetal Life

Constant Supply of Nutrients

- Glucose
- Amino acids
- Lipids
- Calcium, Phosphorus
- Magnesium
- Fluids

Fetal to Neonate Transition

- Neonate must supply and regulate these substances
- Transiently hypoglycemic
 - Low at 2 to 4 hours old
- Hypocalcemia



Nutritional Support

- Early Neonatal Period
- Intravenous fluids
 - For each Kg up to 10 kg - 100 ml/kg/ 24
 - For each Kg from 11-20 kg - 50 ml/kg/ 24
 - For each Kg > 20 kg - 25 ml/kg/ 24 hou
- Dextrose - 4 - 8 mg/kg/min
- Electrolyte concentrations - lab va
 - K – 3 mEq/kg/day
- Until enteral or parenteral nutritio



Early Neonatal Period

Early Hyperglycemia

- Continued glucose production
- Decreased glucose clearance
- Diminished insulin response
- Diminished response
 - Insulin resistance
- Increased cortisol/epinephrine
 - Stress
- Sepsis



Early Neonatal Period

Early Hypoglycemia

- Limited hepatic glycogen stores
- Inadequate endogenous glucose production
 - Failure of transition to gluconeogenesis
- High glucose utilization – hypermetabolism
- High risk for developing hypoglycemia
 - Perinatal asphyxia
 - Placentalitis
 - FIRS
 - Intrauterine growth retardation
 - Placental insufficiency
 - Cold stress
 - Sepsis

Enteral Feeding Advantages

- Physiologic stimulation
- Metabolic regulation
- Gastrointestinal mucosa
- Gastrointestinal mucosa
- Lower cost



Enteral Feeding


Requirements for initiating enteral feeding

- No abdominal distension
- No gastric reflux
- Passage of meconium
- Active bowel sounds
- If severe perinatal distress
 - Stable blood pressure
 - Temperatures near 100 F
 - Normal P_{aO_2}
 - Stable blood glucose





Enteral Feeding Fresh mare's milk

- Preferred source of enteral nutrition
 - Unique nutrient composition
 - Increased bioavailability of nutrients
 - Immunologic properties
 - Promotion of maternal-neonate bonding
 - Present of hormones, enzymes, growth factors
- 
- A photograph of a white mare and her foal in a stable stall. The mare is standing and looking towards the camera, wearing a red and blue halter. The foal is standing next to her, partially obscured. The stall has a concrete block wall and a window with a dark covering. The floor is covered with straw bedding.

Enteral Feeding

Other choices

- Frozen mare's milk
- Goat's milk
- High quality powdered milk replacer



Enteral Feeding

- Initially feed 5% of body weight / 24 hours

Divided into 12 feedings

- If foal tolerates this volume
 - Increase to 10% during the first day
 - Normal foal
 - Target of 20-25% of his body weight / day
- 21% provides the ideal 120 kcal/kg/day

Enteral Feeding

Ill Foal

- Enteral feeding is questionable

Trophic feelings

1-2% body wt divided every 4-6 hrs

Provide calories / protein using parenteral route

- Permissive underfeeding

Over-nutrition associated with sepsis

Ill foals are confined

Target – enough to maintain anabolism

- 1-2% weight gain per day

Usually 12-14% body weight

Enteral Feeding

- Example 20% - 114 kcal/kg/day
- 110 lb Foal
 - 110 lb X 20% = 22 lb
 - 22 lb X 16 oz/lb = 352 oz/day
 - 352 oz / 12 feedings = 29 oz/feeding
- 50 kg Foal
 - 50 kg X 20% = 10 kg = 10 liters
 - 10 liters / 12 feedings = 833 ml/feeding

Enteral Feeding Route

- Suckling
- Small Diameter

Indwelling Nasogastric Tube

Enteral Feeding Suckling

- Best route - physiologic stimulation
- Abnormal suckling behavior
 - Desperately want to suck
 - Ineffective
 - High risk for aspiration
 - Healthy neonate - few consequences
 - Critical neonate - pulmonary disease (sepsis)
- Feed on demand - at least Q2H

Enteral Feeding

Small Diameter Indwelling Nasogastric Tube

- Feed every two hours
- Bolus feeding
- Rhinitis
- Pharyngitis



Parenteral Nutrition

Intravenous delivery of

- Calories
 - Glucose
 - Lipids
- Protein
 - Amino Acids
- Vitamins
- Trace minerals




Parenteral Nutrition

Glucose requirements

- Primary source of energy developing fetus
- Term net umbilical uptake
 - 5 to 8 mg/kg/min (7.2-11.5 gm/kg/day)
- Fetus does not carry out gluconeogenesis

Glucose Support

- All compromise neonates
 - Will benefit from exogenous glucose support
 - Decrease catabolic state
 - Support their recovery
- Blood glucose interpretation
 - Not relate directly to adequacy
 - Summation of glucose mobilization
 - Hypoglycemia
 - Normoglycemia
 - Hyperglycemia



Glucose Measurement

- Bedside monitoring – Glucometers
- Whole blood measurement
 - Electrochemical biosensor
 - Plasma from whole blood
 - Diffuses into, solvates the reagent layer
 - Electrons produced, current generated
 - Photometric test strips
 - Produces a blue color




Glucose Measurement Variation

- PCV
- Total protein
- P_{O_2}
- pH
- Model/Instrument
- Reagent strip
 - Handling
 - Age
 - Lot




Glucose Measurement PCV




Glucose Measurement

PCV and Glucose Level



Glucose Measurement PCV

- Whole blood vs plasma
- Reagent strip sieve plasma
 - Increased hematocrit may mechanically obstruct diffusion
 - Block the “holes” in the mesh membrane
 - Decrease the volume of plasma available to diffuse
 - Alter blood viscosity, decreasing fluid permeability
 - Rouleau formation



Glucose Measurement PCV

- Critically ill neonatal foals
 - Microclot formation
 - Sample hemolysis
 - Protein deposition
 - High fibrinogen levels
 - Fibrin aggregation
 - Platelet/other cellular aggregation
 - Other inflammatory phenomena



Parenteral Nutrition

Glucose requirements

- At birth glucose
 - Gluconeogenesis (catecholamine secretions)
 - Hepatic glycogenolysis
 - Umbilical cord rupture - release Glucagon
- At birth the stimulated fetal liver
 - 4 to 8 mg glucose/kg/minute

Parenteral Nutrition Protein Requirements

- Estimate amino acid utilization late term fetus
- Intrauterine nitrogen delivering - lambs
2.7 to 3.5 gm/kg/day
When total energy is > 70 kcal/kg/day
- Increased requirements
 - Stress
 - Infection
 - SIRS

Parenteral Nutrition

Lipid requirements

- Lipids are not utilized by fetus as energy
- Periods of stress - very important
- Neonatal foals utilize lipids as an energy source

40 - 50% of total caloric intake can come from fat

Parenteral Nutrition

Starting Formula

Example - 50 kg foal

- Dextrose - 10 gm/kg/day - 34 kcal/kg
- Amino acids - 2 gm/kg/day - 8 kcal/kg
- Lipids - 1 gm/kg/day - 11 kcal/kg
- Plus vitamins and trace minerals
- Total - 53 kcal/kg
- Nonprotein kcal - 24% lipid
- Nonprotein kcals:gram nitrogen = 139.5

Foal Nutrition

Numbers to Know

- Glucose - 3.4 kcal/gm
- Amino acids - 4.0 kcal/gm
- Lipid - 9.0 kcal/gm
- 6.25 gm amino acids = 1 gm nitrogen
- 1 oz = 30 ml
- 16 oz milk = 1 lb
- Mare's milk - 0.57 kcal/ml

Maternal Bonding Behavioral Therapy

- Mares need contact with foals
 - Sight
 - Smell
 - Touch
 - Hear
- Stimulus for lactation
 - Establish milk flow
 - Maintain milk flow



Nurse Mare Bonding

- Environment
 - Quiet
 - No distractions
 - "Foal's Stall"
- Foal's behavior
 - Vigorous nursing
 - Aggressive nursing
- Foal's Smell
- Foal's physical appearance

